

Rx Only 
NICADAN[®]

Niacinamide, Vitamin B6, Vitamin C, Folic Acid,
Copper, Magnesium, Zinc & Alpha Lipoic Acid

DESCRIPTION

Nicadan[®] tablets are a specially formulated dietary supplement containing natural ingredients with anti-inflammatory properties. Each pink-coated tablet is oval shaped, scored and embossed with "MM". **Nicadan**[®] is for oral administration only.

Nicadan[®] should be administered under the supervision of a licensed medical practitioner.

INGREDIENTS

Each tablet of **Nicadan**[®] contains:

Vitamin C (as Ascorbic Acid)	100 mg
Niacinamide (Vitamin B-3)	800 mg
Vitamin B-6 (as Pyridoxine HCl)	10 mg
Folic Acid	500 mcg
Magnesium (as Magnesium Citrate)	5 mg
Zinc (as Zinc Gluconate)	20 mg
Copper (as Copper Gluconate)	2 mg
Alpha Lipoic Acid	50 mg

Other Ingredients: Microcrystalline cellulose, Povidone, Hypromellose, Croscarmellose Sodium, Polydextrose, Talc, Magnesium Sterate Vegetable, Vegetable Stearine, Red Beut Powder, Titanium Dioxide, Maltodextrin and Triglycerides.

Nicadan[®] is free of yeast, wheat, dairy, egg, soy, gluten, artificial color or flavors, added sugar, starch and preservatives.

USES

As a dietary supplement, **Nicadan**[®] tablets are beneficial in maintaining overall skin health, especially in patients with inflammatory skin conditions.*

**This statement has not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure or prevent any disease.*

WARNINGS

If you are pregnant, nursing or taking medication, consult your doctor before use.

CONTRAINDICATIONS

Use of **Nicadan**[®] is contraindicated in patients who are hypersensitive to any ingredient in this tablet.

PRECAUTIONS

There have been reports of GI upset, nausea, vomiting, heart-burn, diarrhea, dizziness, drowsiness or headache with large doses of niacinamide. Rarely, niacinamide in doses of 3 grams/day or more has been associated with elevated liver function tests. There are case reports in the literature indicating that

niacinamide may reduce the hepatic metabolism of primidone and carbamazepine. Individuals taking these medications should consult their physician. Individuals taking anti-diabetes medications should have their blood glucose levels monitored.

Allergic sensitization has been reported rarely following oral administration of folic acid. Folic acid above 1 mg daily may obscure pernicious anemia in that hematologic remission may occur while neurological manifestations remain progressive.

DOSAGE AND ADMINISTRATION

Take one tablet daily with food or as directed by a physician.

Nicadan[®] tablets are scored, so they may be broken in half if required.

HOW SUPPLIED

Nicadan[®] is available in a bottle containing 60 tablets. 43538-440-60

Store at 15°C to 30°C (59°F to 86°F). Keep bottle tightly closed. Store in cool dry place.

KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.

You may report side effects by calling 1-973-882-7512.

Manufactured for:



383 Route 46 West, Fairfield, NJ 07004-2402 USA

www.medimetriks.com

Rev.: 5/2015